

Herbal Medicines Compendium

Published on *Herbal Medicines Compendium* (<https://hmc.usp.org>)

Panax ginseng Steamed Root and Rhizome Dry Extract

Proposed For Development Version 0.1

Panax ginseng Steamed Root and Rhizome Dry Extract

DEFINITION

The article is prepared from the dried root and rhizome of cultivated *Panax ginseng* C.A.Mey. (Family Araliaceae) collected in the fall, steamed until the material externally reddish-brown and translucent, by extraction with water or hydroalcoholic mixtures. It contains NLT 90.0% and NMT 110.0% of the labeled amount of total ginsenosides, calculated as the sum of ginsenoside Rg1 ($C_{42}H_{72}O_{14}$), ginsenoside Re ($C_{48}H_{82}O_{18}$), ginsenoside Rf ($C_{42}H_{72}O_{14}$), ginsenoside Rb1 ($C_{54}H_{92}O_{23}$), ginsenoside Ro ($C_{48}H_{76}O_{19}$), ginsenoside Rc ($C_{53}H_{90}O_{22}$), ginsenoside Rb2 ($C_{53}H_{90}O_{22}$), and ginsenoside...

Source URL (modified on 2016/02/17 - 3:11pm):<https://hmc.usp.org/monographs/panax-ginseng-steamed-root-and-rhizome-dry-extract-0-1>